## Amendments to the Claims:

This listing of clams will replace all prior versions, and listing, of claims in the application:

## **Listing of Claims:**

1. (Original): A protein binding assay for measuring IP<sub>3</sub> in a sample employing as reagents a conjugate of IP<sub>3</sub> and a detectable label joined through a bond or linker at the 2-hydroxyl position, and a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R, said method comprising:

combining in an assay medium said sample, said conjugate and said binding protein and incubating said mixture for sufficient time for any IP<sub>3</sub> and said conjugate to bind to said binding protein; and

detecting the bound or unbound label as a measure of the IP<sub>3</sub> present in the sample.

- 2. (Original): A protein binding assay according to Claim 1, wherein said assay is in a homogeneous format.
- 3. (Original): A protein binding assay according to Claim 1, wherein said sample is a cellular lysate, and wherein said cellular lysate has been treated to block kinases and phosphatases and prepare said sample for said assay.
- 4. (Original): A protein binding assay according to Claim 1, wherein said binding protein is of not more than about 600 amino acids and comprises at least amino acids 226 578 of the mouse IP<sub>3</sub>R Type 1.
- 5. (Original): A protein binding assay according to Claim 1, wherein said label is an enzyme fragment for enzyme complementation.

- 6. (Original): A protein binding assay according to Claim 1, wherein said binding protein is a fusion protein of up to about 1.5kD amino acids.
- 7. (Original): A protein binding assay according to Claim 1, wherein said label is a fluorescer.
- 8. (Original): A method according to Claim 1, wherein the order of addition of reagents is: (a) combining said sample with said binding protein; and (b) adding said conjugate, with incubating after (a) and (b).
- 9. (Original): A protein binding assay for measuring IP<sub>3</sub> in a sample using a homogeneous format, employing as reagents a conjugate of IP<sub>3</sub> and an ED of from 37 to 60 amino acids derived from β-galactosidase joined through a linker at the 2-hydroxyl position, and a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R, said method comprising:

combining in an assay medium assay components in the following order: said sample, said binding protein, said conjugate and EA, and incubating after each combining for sufficient time for complex formation between said assay components;

adding substrate for said  $\beta$ -galactosidase; and

detecting the turnover of said  $\beta$ -galactosidase of said substrate as a measure of the IP<sub>3</sub> present in the sample.

10. (Original): A protein binding assay for measuring IP<sub>3</sub> in a sample using a homogeneous format, employing as reagents a conjugate of IP<sub>3</sub> and a fluorescer joined through a linker at the 2-hydroxyl position, and a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R, said method comprising:

combining in an assay medium assay components: said sample, said binding protein, and said conjugate, and incubating for sufficient time for complex formation between said assay components; and

detecting the change in fluorescence polarization as a measure of the IP<sub>3</sub> present in the sample.

- 11. (Original): A method according to Claim 10, wherein said linker is an aliphatic group of from 4 to 20 carbon atoms.
- 12.(Currently amended): A method according to Claim 9 Claim 10, wherein said fluorescer emits at a wavelength greater than about 500 nm.
- 13. (Original): A method according to Claim 10, wherein said fluorescer has a polarizability of less than about 60mP.
- 14. (Original): A protein binding assay for measuring IP<sub>3</sub> in a sample employing as reagents a conjugate of IP<sub>3</sub> and a detectable label joined through a bond or linker at the 2-hydroxyl position, and a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R, said method comprising:

combining in an assay medium said sample, said conjugate, said binding protein and a chemical reductant and incubating said mixture for sufficient time for any IP<sub>3</sub> and said conjugate to bind to said binding protein; and

detecting the bound or unbound label as a measure of the IP<sub>3</sub> present in the sample.

15. (Original): A protein binding assay according to Claim 14, wherein said chemical reductant is a thiol.

Attorney Docket No. 3817.14-1 Customer No. 23308

## 16. (Original): A compound of the formula:

wherein:

R is a neutral linking group of from 4 to 20 carbon atoms bonded to the oxygen through a saturated carbon atom or carbonyl;

Z is a functionality for linking X to the oxygen at the 2-position;

X is an enzyme donor fragment of  $\beta$ -galactosidase of from 27 to 60 amino acids; and

n is 1 or 2.

17. (Original): A compound of the formula:

Attorney Docket No. 3817.14-1 Customer No. 23308

wherein:

R is a neutral linking group of from 2 to 20 carbon atoms bonded to the oxygen through a saturated carbon atom;

Z is a functionality for linking X to the oxygen at the 2-position; and X is a fluorescer.

- 18. (Original): A kit comprising a compound according to Claim 17, enzyme acceptor for said enzyme donor and a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R.
- 19. (Original): A kit comprising a compound according to Claim 18, enzyme acceptor for said enzyme donor and a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R.
- 20. (Original): A kit for performing an IP3 assay comprising a conjugate of IP<sub>3</sub> and a detectable label joined through a bond or linker at the 2-hydroxyl position, a truncated extracellular portion of an IP<sub>3</sub>R having at least about 200 times the affinity for IP<sub>3</sub> than the intact IP<sub>3</sub>R and instructions for performing said assay.

Attorney Docket No. 3817.14-1 Customer No. 23308

21. (Original): A kit according to Claim 20, further comprising a thiol reductant.